

# A Novel Thermostable GH51 $\alpha$ -L-Arabinofuranosidase From *Caldicellulosiruptor Lactoaceticus* and The Application for Natural Hemicellulose Degradation

Xiaojing Jia<sup>1,2,3</sup>, Meng Li<sup>1</sup>, Yejun Han<sup>2\*</sup>

<sup>1</sup>Beijing Advanced Innovation Center for Food Nutrition and Human Health, Beijing Technology and Business University

<sup>2</sup>National Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100190, China

<sup>3</sup>University of Chinese Academy of Sciences, Beijing 100049, China

Volume 1 Issue 1- 2018

Received Date: 26 July 2018

Accepted Date: 30 Aug 2018

Published Date: 05 Sep 2018

## 2. Keywords

$\alpha$ -L-Arabinofuranosidase; Pentoses production; *Caldicellulosiruptor lactoaceticus*; Arabinoxylan; Synergy; Thermophilic degradation

## 1. Abstract

Improving the hydrolytic performance of xylanolytic enzymes on complicated xylan is of great importance in lignocellulosic biomass-based fermentation industry. Employing a de-branching enzyme  $\alpha$ -L-arabinofuranosidase (EC 3.2.1.55) in conjunction with backbone depolymerizing enzyme endo- $\beta$ -1,4-xylanase (EC 3.2.1.8) offers a way to address this problem. In present study, a glycoside hydrolase family 51  $\alpha$ -L-arabinofuranosidase Abf51A from extremely thermophilic bacterium *Caldicellulosiruptor lactoaceticus* was cloned, expressed and characterized in *Escherichia coli*. The recombinant Abf51A existed as a homohexamer in solution, which showed optimal activity at 80°C and pH 5.5, and had long-term thermostability at 75°C and 80°C after 12 h incubation. Abf51A was most active on p-nitrophenyl- $\alpha$ -L-arabinofuranoside with  $K_m$  and  $V_{max}$  values reaching  $0.52 \pm 0.05$  mM and  $2981.00 \pm 79.65$   $\mu$ mol/min/mg, respectively. In addition, it also displayed minor activity with p-nitrophenyl- $\beta$ -D-xylopyranoside. Polymeric substrates screening revealed that Abf51A released arabinose from arabinoxylan (oat spelt xylan) and glucuronoarabinoxylan (sugarcane xylan and corn cob xylan). Moreover, Abf51A showed obvious synergism with endo- $\beta$ -1,4-xylanase Xyn10A on both arabinoxylan and glucuronoarabinoxylan degradation, suggesting a significant contribution of Abf51A in arabinose-substituted xylan utilization. The thermostability, hydrolytic properties and synergism action made Abf51A a promising candidate in the production of fermentable pentoses from arabinose-containing xylylans.

## 3. Introduction

Xylan, as the most abundant hemicellulosic constituent of plant cell walls, has been considered as an abundant source of fermentable pentose (mainly xylose and arabinose) for biofuels and biochemicals. Generally, it consisted of a linear backbone of  $\beta$ -1,4-linked D-xylopyranosyl units which can be degraded by core enzymes endo- $\beta$ -1,4-xylanase (EC 3.2.1.8) and  $\beta$ -xylosidase (EC 3.2.1.37) (Dodd and Cann 2009). Xylan is usually linked with various side-chain substitutions to its main chain, such as L-arabinofuranosyl, acetyl, feruloyl and glucuronic acids residues. Hence xylylans are grouped into linear homoxylan, arabinoxylan (AX), glucuron-

oxylan, glucuronoarabinoxylan (GAX) and heteroxylan based on these substituents on the backbone [1]. However, the presence of these side chains on xylan can strongly restrict the action of endo- $\beta$ -1,4-xylanase and  $\beta$ -xylosidase, thus preventing the complete breakdown of the polymer to its basic units [2]. Therefore, among the xylanolytic enzyme system, de-branching enzymes such as  $\alpha$ -L-arabinofuranosidase (EC 3.2.1.55, AFase),  $\alpha$ -glucuronidase (EC 3.2.1.131) and acetyl xylan esterase (EC 3.1.1.72) are required for the synergistically enzymatic action on polymeric xylan. Currently, these side chain-removing accessory enzymes have been used collaboratively with other lignocellulose-decomposing en-

\*Corresponding Author (s): Yejun Han, National Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing, China, Email: yejunhan09@gmail.com

zymes in biotechnological processes [3]. AFases are hemicellulases that cleave the 1-2, 1-3 and 1-5 glycosidic bond at the non-reducing terminal ends between L-arabinofuranoside substituents and various poly- and oligo-saccharides. These enzymes are part of glycoside hydrolases (GHs) responsible for the degradation of arabinose-existing hemicelluloses such as L-arabinan, AX and arabinogalactan [4]. Based on the amino acid sequence similarities and hydrophobic cluster analysis, AFases have been classified into GH families 3, 43, 51, 54 and 62, with families GH51, 43 and 62 as the main families (CAZY; <http://www.cazy.org/>). To date, GH51 is the largest characterized family with a majority of them being of bacterial origin. Structure data obtained for several members of this family confirm that GH51 AFases process a catalytic domain with  $(\beta/\alpha)_8$ -barrel architecture, and additionally uptake a unknown function domain composed of  $\beta$ -sandwich at the C-terminal. Besides, GH51 enzymes preform glycosidic bond cleavage *via* a retaining mechanism with two conserved glutamate residues acting as acid/base and nucleophile. So far, the majority of characterized GH51 AFases showed only exo-arabinanase activity with arabino-oligosaccharides and/or L-arabinose containing heteropolymers such as arabinan and AX [5]. A few GH51 AFases also display endo- $\beta$ -D-xylanase activity [6]. A finely planned cocktail of enzymes, containing not only cellulases but also xylanases, is required for efficient fermentable sugars yield from the polysaccharide components of lignocellulosic biomass. In this respect, thermostable enzymes process a number of important advantages compared with their mesophilic counterparts in the optimization of biorefinery-scale lignocellulosic depolymerization [7]. Various bacteria, archaea, fungi and plants have received attentions as sources for thermostable xylanolytic enzymes. Thermophilic microbes provide convenient approaches to obtain thermostable enzymes active on lignocellulose. *Caldicellulosiruptor lactoaceticus* 6A, an extremely thermophilic plant biomass degrading bacterium, grows on natural xylan as sole carbon source [8]. In our previous studies on glycoside hydrolases for de-branched xylan degradation from this bacterium, a thermophilic endo- $\beta$ -1,4-xylanase Xyn10A with good thermostability has been reported [9]. Further insight into xylanolytic enzymes other than endo- $\beta$ -1,4-xylanase of this bacterium would provide not only new exploration to the entire mechanisms for xylan utilization of *C. lactoaceticus*, but also approach for biomass conversion at extreme temperature. In present study, the gene encoding AFase Abf51A was identified and cloned from the genome of *C. lactoaceticus* 6A, then expressed in *E. coli* and biochemically characterized. Additionally, the synergistically hydrolytic function of AFase together with endo- $\beta$ -1,4-xylanase on heteroxylan hydrolysis was also studied.

## 4. Materials and Methods

### 4.1. Strains, plasmids and chemicals

*C. lactoaceticus* 6A (= DSM 9545) was obtained from DSMZ (Braunschweig, Germany). *Escherichia coli* Top10 (TianGen, China) and plasmid pET-28b (Novagen, USA) were used for gene cloning, and *E. coli* Rosetta (DE3) (Novagen, USA) was used for protein expression and purification. *p*-nitrophenyl- $\alpha$ -L-arabinofuranoside (*p*NP-Af), *p*-nitrophenyl acetate (*p*NP-Ace), *p*-nitrophenyl- $\alpha$ -D-xylopyranoside (*p*NP- $\alpha$ -Xyl), *p*-nitrophenyl- $\beta$ -D-xylopyranoside (*p*NP- $\beta$ -Xyl), *p*-nitrophenyl- $\beta$ -D-cellobioside (*p*NP-C) and *p*-nitrophenyl- $\beta$ -D-glucopyranoside (*p*NP-Glu) were purchased from Sigma-Aldrich (St. Luis, USA). Oat spelt xylan (OSX) was purchased from Hualan Chemical Technology Co., Ltd. (Shanghai, China). Corn cob xylan (CCX) and sugarcane xylan (SCX) was presented by Professor Hourui Zhang from Guangxi Institute of Botany, Chinese Academy of Sciences. D-xylose, L-arabinose, locust bean gum, soluble starch, Avicel, carboxymethyl cellulose (CMC), *p*-nitrophenol, and other chemicals were obtained from Kepujia Reagent Co. (Beijing, China). All other chemicals were of analytical class unless otherwise stated.

### 4.2. Genomic DNA extraction and amplification

*C. lactoaceticus* 6A was cultivated in DSM 671 medium at 70°C for 4 d under anaerobic conditions. The genomic DNA of *C. lactoaceticus* 6A was isolated using a TIANamp Bacteria DNA Kit (TianGen, Beijing, China). The gene *abf51A* encoding a hypothetical AFase [GenBank: AEM73175.1] was predicted and amplified based on the whole genome [10]. PCR reaction was performed with Pfu DNA polymerase (TianGen) and the procedure was as follows: 95°C 5 min; 30 cycles of 95°C 30 s, 55°C 1 min, and 72°C 1 min; final 72°C 5 min. Target PCR products were then purified using TIAN Gel Midi Purification Kit (TianGen) and stored at -20°C until use.

### 4.3 Construction and sequencing of the expression vector

The purified PCR products of *abf51A* amplification were then digested with T4 DNA polymerase (Takara, Dalian, China) and inserted into pET-28b EK/LIC vector. The sub-cloned plasmid was then transferred into *E. coli* Top10 competent cells by heat shock and grown overnight on Lysogeny Broth (LB) agar plate containing 50  $\mu$ g/mL kanamycin at 37°C. Positive cloning junctions were confirmed by colony PCR and DNA sequencing with T7 primers from both strands.

### 4.4. Expression, purification and gel filtration chromatography

The recombinant plasmid was isolated using TIANprep Mini Plasmid Kit (TianGen) and transferred into *E. coli* Rosetta (DE3)

competent cells by heat shock and grown overnight on LB agar plates containing 50 µg/mL kanamycin and 34 µg/mL chloramphenicol at 37°C. One single colony was picked out as seed culture and grown overnight in 5 mL LB liquid medium containing 50 µg/mL kanamycin and 34 µg/mL chloramphenicol at 37°C on a rotary shaker (220 rpm). The culture was induced with 0.5 mM Isopropyl-β-D-thiogalactopyranoside (IPTG) after the optical density reached 0.6 at 600 nm for an additional 6 h at 37°C. The cells were harvested by centrifugation (4000 rpm for 20 min at 4°C), and resuspended in 20 mL binding buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl). Then the cell suspension was lysed by sonication on ice and cell debris was removed by centrifugation (12000 rpm for 20 min at 4°C). After then, the supernatant was heat-treated at 50°C for 30 min and denatured proteins were removed by centrifugation (12000 rpm for 20 min at 4°C). The resulting supernatant was then loaded onto a Ni-NTA affinity column (National Engineering Research Centre for Biotechnology (NERCB), Beijing, China), which was pre-equilibrated with binding buffer for five times. Subsequently, the column was washed with binding buffer for three times and eluted with elution buffer (50 mM Tris-HCl, pH 7.5, 300 mM NaCl, 150 mM Imidazole). Hereafter, the molecular mass and oligomeric state were further determined using a 1 × 33 cm<sup>3</sup> Superdex 200 exclusion column (Mengyimei Bio-Tech Co., Ltd. Beijing, China) using citrate buffer (50 mM sodium citrate, 150 mM NaCl, pH 6.0) as the mobile phase. Approximate 500 µL sample was loaded onto the column at a flow rate of 0.5 mL/min using Huxi chromatographic separation system (Shanghai Huxi analysis instrument factory, Co. LTD, China). At the same time, the homogeneity of the target protein was checked by sodium dodecyl sulfate-polycrylamide gel electrophoresis (SDS-PAGE) in accordance with the technique described by Laemmli [11].

#### 4.5. Enzyme assay and protein determination

The activity of AFase was measured using *p*NP-Af as the substrate as described by Qiaojuan Yan [12] with slight modifications. The reaction mixture, contained diluted enzyme, 1 mM *p*NP-Af and 50 mM citrate buffer (pH 5.5) in a 50 µL reaction volume, was incubated at 80°C for 3 min. The reaction was then stopped by adding 100 µL of 1 M Na<sub>2</sub>CO<sub>3</sub> and the number of *p*-nitrophenol generated was measured at 400 nm. A standard curve was prepared with *p*-nitrophenol as a standard. One unit of enzyme activity was defined as the amount of enzyme required to release 1 µmol of *p*-nitrophenol per minute under the standard assay conditions. Protein concentration was determined by the Bradford method using bovine serum albumin (BSA) as the standard. All of the assays were performed in triplicate.

#### 4.6. Biochemical characterization of the recombinant enzyme

Rapid screening of Abf51A activity was firstly conducted with various aryl-monosaccharides including *p*NP-Af, *p*NP-Ace, *p*NP-β-Xyl, *p*NP-α-Xyl, *p*NP-C and *p*NP-Glu. Besides, the substrate specificity was also detected with 1% (*w/v*) polysaccharides including OSX, CCX, SCX, locust bean gum, soluble starch, Avicel and CMC. One unit of Abf51A activity towards polysaccharides was defined as the amount of enzyme required to release 1 µmol of L-arabinose equivalent per minute under the standard assay conditions. The optimum temperature for Abf51A activity was measured in a temperature range of 40-95°C and pH 6.5 with *p*NP-Af as substrate as described above. In addition, the optimum pH for Abf51A activity was determined by using 50 mM phosphate-citrate buffers (pH 4.0-8.5) at 75°C with *p*NP-Af as substrate. To detect the thermostability of the enzyme, purified Abf51A were incubated in 50 mM phosphate-citrate buffer (pH 6.0) at 75, 80 and 85°C for different time periods in the absence of substrate. The residual activity of Abf51A under optimal conditions was assayed at the time period of 0.5, 1, 2, 4, 6 and 12 h with *p*NP-Af as substrate. All of the experiments were performed in triplicate. Moreover, the kinetic parameters  $K_m$  and  $V_{max}$  for Abf51A were determined by incubating enzyme with *p*NP-Af at various concentrations (0.01-5.0 mM) under optimal conditions and calculated from a Lineweaver-Burk plot using GraphPad Prism version 6.0 for Windows (GraphPad Software Inc., USA).

#### 4.7. Synergistic hydrolysis of xylan

To investigate the synergistic effect of Abf51A in combination with β-1,4-xylanase, the GH family 10 xylanase Xyn10A from *C. lactoaceticus* 6A was purified as previously described [13]. The reaction mixture (1 mL) contained 1% (*w/v*) substrates (OSX, CCX and SCX), 50 mM citrate buffer (pH 6.0), and 25 µg purified enzyme (Abf51A and Xyn10A, either alone or in combination). Each of the control reaction was performed under the same experimental condition except adding corresponding heat-denatured enzymes. After incubation at 80°C for 24 h, the reaction was stopped by boiling for 10 min, and centrifuged at 12000 rpm for 10 min. The amount of reducing sugars was determined using *para*-hydroxybenzoic acid hydrazide (PHBAH) method with D-xylose as a standard [14]. At the same time, the released D-xylose and L-arabinose were determined by high performance liquid chromatography (HPLC) using a Hi-Plex Ca column (300 × 7.7 mm, Agilent Technologies, United Kingdom) and a refractive index detector (LC-20AT, Shimadzu Corp., Japan). The column temperature was maintained at 85°C with Milli-Q filtered HPLC grade water as mobile phase at a flow rate of 0.6 mL/min. The injection volume was 10 µL with D-xylose and L-arabinose as standards. All the assays were performed for three times.



5. Results

5.1. Sequence analysis and phylogenetic analysis

Through *C. lactoaceticus* genome sequence analysis, Calla\_0521 was annotated as a putative AFase (Abf51A). Abf51A was predicted to have no signal peptide indicating it was an intracellular enzyme. The *abf51A* gene was amplified to encode 505 amino acid residues, and the calculated molecular weight (Mw) and deduced theoretical isoelectric point (pI) of the protein were 57.87 kDa and 5.83, respectively. Abf51A showed the highest (99%) identity with AFases from *Caldicellulosiruptor acetigenus* [GenBank: WP\_029228247.1] and *Caldicellulosiruptor kristjanssonii* [GenBank: WP\_013432406.1], and higher (92-98%) identity with other predicted AFases from *Caldicellulosiruptor* sp. Besides, Abf51A exhibited 70% identity with AFase from thermophilic bacteria *Dictyoglomusthermophilum* [GenBank: WP\_012548731.1], *Dictyoglomusturgidum* [GenBank: WP\_012584082.1] and *Mahellaaustraliensis* [GenBank: WP\_013779842.1]. Moreover, to gain further insight into the evolutionary relationship between GH51 AFases, phylogenetic tree was constructed using 18 protein sequences of GH51 AFases by Neighbor-Joining method (Figure. 1). The results revealed that Abf51A was clustered with GH51 AFase from *Caldicellulosiruptor saccharolyticus* DSM 8903 [GenBank: ABP67153.1], then located closer with GH51 AFases from *Paenibacillus* sp. Besides, amino acids sequence blast analysis revealed that Abf51A contained an Alpha-L-AF-C superfamily motif (291-494 amino acids) that had significant similarity with other GH51 AFases. Aligning the Abf51A sequence with several GH51 AFases revealed that they shared the conserved catalytic residues, that was Glu173 acting as the acid/base, and Glu292 acting as the catalytic nucleophile (Figure. 2).






Figure 2: Multiple amino acid sequence alignment of Abf51A. Sequence alignment was computed on <http://www.genome.jp/tools/clustalw/> and depicted by ESPrnt 3.0: <http://esprnt.ibcp.fr/ESPrnt/cgi-bin/ESPrnt.cgi>, respectively. Identical and similar amino acid residues were enclosed in black boxes and open boxes, respectively. The catalytic residues Glu173 and Glu292 of Abf51A were marked by black stars below the conserved sequences. The alignment for Abf51A included GH51 AFases from *Geobacillus stearothermophilus* T6 [PDB: 1PZ3\_A], *Ruminiclostridium thermocellum* [PDB: 2C7F\_A], *Bifidobacterium longum* [PDB: 2Y2W\_A], *Thermotoga petrophila* RKU-1 [PDB: 3S2C\_A] and *Thermotoga maritima* MSB8 [PDB: 4ATW\_A].

Figure 1: Phylogenetic tree analysis of Abf51A. The Neighbor-Joining tree was constructed using MEGA 5.05, bootstrap value was 1000 replicates, and GenBank accession numbers of each protein sequence were listed after scientific names.

## 5.2. Structure analysis


The structure of Abf51A was built by homology modeling from Swiss Model server (Figure. 3). The closest monomer homolog was GH51 AFase from *G. stearothermophilus* T6 [PDB: 1PZ3\_A], with which Abf51A shares 65.1% identity and for which the structure-function relationship was defined. The overall two domain structure of Abf51A monomeric subunit composed of two domains: a  $(\beta/\alpha)_8$  barrel and a C-terminal 12-stranded  $\beta$ -sandwich with a jelly-roll topology. The homology model of Abf51A also confirmed the putative role of acid/base catalyst Glu173 and nucleophile residue Glu292.



**Figure 3:** Structural model of Abf51A. Structure was created by Swiss-model and processed by Chimera 1.6.2. The structure of Abf51A was built with the GH51 AFase from *G. stearothermophilus* T6 [PDB: 1PZ3\_A] as template. The catalytic residues were shown as yellow sticks.

## 5.3. Purification and gel filtration chromatography property

For functional analysis of the recombinant enzyme, Abf51A was cloned into pET-28b vector and expressed in *E. coli* Rosetta (DE3). The recombinant protein from the crude extract was purified by Ni-affinity chromatography and further determined through Superdex 200 gel filtration. Recombinant Abf51A displayed as a single band with an Mw of about 57 kDa after SDS-PAGE analysis, which was consistent with the predicted Mw based on the amino acid sequence (Figure.4a). Size exclusion chromatography revealed that Abf51A eluted as a single peak corresponding to Mw of 340 kDa, suggesting that Abf51 existed as a homohexameric solution (Figure.4b).



**Figure 4:** SDS-PAGE analysis (a) and gel filtration chromatography profile (b) of Abf51A.

## 5.4. Biochemical characterization of the recombinant enzyme

Rapid screening of Abf51A activity using different aryl-monosaccharides confirmed that Abf51A was only specific for *p*NP-Af (2770.83 U/mg) and *p*NP- $\beta$ -Xyl (2.32 U/mg), but not for other aryl-glycosides including *p*NP-Ace, *p*NP- $\alpha$ -Xyl, *p*NP-C and *p*NP-Glu. Further testing on polymeric substrates revealed that Abf51A could release trace amount of reducing sugars from a variety of xylan polymers including OSX, CCX and SCX with specific activity of 0.006, 0.003 and 0.001 U/mg, respectively. Besides, no activities were observed for locust bean gum, soluble starch, Avicel and CMC. These results indicated that the recombinant Abf51A was an exo-type arabinanase. The Michaelis-Menten constant  $K_m$  and  $V_{max}$  values for *p*NP-Af under standard assay conditions were  $0.52 \pm 0.05$  mM and  $2981.00 \pm 79.65$   $\mu$ mol/min/mg, respectively.

Analysis of enzymatic activity revealed that optimal temperature and pH for Abf51A were 80°C and 5.5, respectively. Abf51A retained more than 55% of its activity at 60-85°C, and maintained over 60% of the activity at pH 4.5-6.5, while decreased rapidly of more than 60% of its activity at pH 7.0-8.5, suggesting its thermophilic and acidic activity in solution (Figure.5a). Moreover, Abf51A was incubated at 75, 80 and 85°C in the absence of substrate for thermostability determination (Figure.5b). Abf51A withstood long-term incubation at 75°C, retaining 63.0% of its original activity after 12 h at this temperature. Even at 80°C, 56.9% of the residual activity was left after 12 h, while 57.0% of the enzyme's activity dropped sharply at 85°C after 30 min.


Moreover, several properties of Abf51A were compared with some other thermophilic AFases from bacteria and fungi as shown in Table 1. The activity of thermostable AFases was maximal at pH 5.0-7.0. Abf51A was evidently distinguished from other AFases with much higher  $V_{max}$  value for *p*NP-Af. Besides, together with Abf51A, some AFases also had scant activity towards *p*NP- $\beta$ -Xyl and xylan polymers.

## 5.5. Synergy with endo- $\beta$ -1,4-xylanase


The applications of AFase in combination with endo- $\beta$ -1,4-xylanase was studied in the hydrolysis of arabinose-rich xylans (Figure.6). Under the tested conditions (80°C, pH 6.0 for 24 h), both Abf51 and Xyn10A gave optimal activities, and remained more than 55% of their original activities after prolonged incubation. The pronounced release of reducing sugars from all tested xylans were achieved in the presence of Xyn10A compared with the separate use of Abf51A, but maximal substrate hydrolysis was attained in the combination of the two enzymes (Figure. 6a). During the hydrolytic process, a large amount of xylose was produced by Xyn10A alone since xylan was mainly consisted of  $\beta$ -1,4-linked D-xylopyranosyl backbone. While much more xylose was released by the simulta-



neous action of Xyn10A and Abf51A, resulting in the highest 2.46-fold increase over the action of Xyn10A alone towards OSX (Figure. 6b). At the same time, Abf51A individually liberated the highest number of arabinose from OSX, but led to the biggest enhancement of arabinose production by 2.25 fold together with Xyn10A on CCX (Figure. 6c). The most significant synergy between Xyn10A and Abf51A was obtained with OSX as substrate compared with that of CCX and SCX (Figure. 6d).



**Figure 5:** Effects of temperature and pH on the activity and stability of the recombinant Abf51A. (a) Temperature and pH optimum. (b) Temperature stability. Optimum temperature was measured at temperatures ranging from 40 to 95°C in phosphate-citrate buffer (pH 6.5). Similarly, optimum pH was determined in phosphate-citrate buffers (pH 4.0-8.5) at 75°C. For investigating thermostability, purified proteins were respectively incubated at 75, 80 and 85°C for different time periods in the absence of substrate. The residual activity was assayed under optimal conditions with pNP-Af as substrate. The highest activity was defined as 100% in triplicate.



**Figure 6:** Synergistic hydrolysis of arabinose-rich xylans. (a) Reducing sugars assay. (b) Released xylose from xylans. (c) Released arabinose from xylans. (d) HPLC analysis of the hydrolytic products of OSX. The reactions were conducted in 1 mL mixture (pH 6.0) containing 25 µg purified protein (Abf51A and Xyn10A, either alone or in combination) and 1% (w/v) xylans (OSX, CCX and SCX) at 80°C for 24 h. The results represented the average of three experimental repeats.

Microorganism	Mw <sup>a</sup> (kDa)	Topt (°C)	pHopt	V <sub>max</sub> <sup>b</sup> (µmol/min/mg)	K <sub>m</sub> <sup>b</sup> (mM)	Spe act <sup>c</sup> (U/mg)	Xylan polymer attacked	References
<i>C. lactoaceticus</i>	57 (340)	80	5.5	2981.00	0.52	2.32	OSX, CCX, SCX	This study
<i>Anoxybacillus thermoautotrophicus</i>	58 (230)	65	5.5	1019.00	0.14	0	SBA, AX, OSX	Canakci et al. 2008
<i>Geobacillus caldophilus</i> TK4	59 (236)	75-80	6.0	588.20	0.17	151.50 <sup>d</sup>	NA	Canakci et al. 2007
<i>Thermobacillus xylanolyticus</i>	56 (NA)	75	5.6-6.2	555.00	0.50	NA	AX, LWX, OSX	Debeche et al. 2000
<i>Paenibacillus</i> sp. DG-22	57 (222)	60	6.0	306.10	3.50	0.02	NA	Lee and Lee 2014
<i>Ruminococcus</i> 7	56 (NA)	50	6.0	270.30	1.05	< 0.01	BeeWX, WFA	Yang et al. 2016
<i>Caldicellulosiruptor saccharolyticus</i>	58 (460)	80	5.5	186.00	1.29	0	NA	Lim et al. 2010
<i>Thermotoga thermarum</i> DSM5069	55 (NA)	95	5.0	162.97	0.19	NA	NA	Xie et al. 2016
<i>Paenibacillus</i> sp. HanTMS1	57 (NA)	75	5.0	NA	0.31	1.70	BirWX, BeeWX	Bouroufi et al. 2016
<i>Thermotoga maritima</i> MSB8	55 (332)	90	7.0	NA	0.42	2.02 <sup>d</sup>	NA	Miyazaki 2005
<i>Clostridium stercorarium</i>	52 (195)	70	5.0	NA	NA	< 0.01	OSX	Schwarz et al. 1995
<i>Geobacillus</i> sp. KCTC 3012	65 (NA)	60	6.0	NA	NA	NA	OSX	Arti et al. 2012

\*Abbreviations: Mw, molecular weight; Topt, optimum temperature; pHopt, optimum pH; Spe act, Specific activity; NA, not available; OSX, oat spelt xylan; CCX, corncob xylan; SCX, sugarcane xylan; SBA, sugar beet arabinan; AX, arabinoxylan; LWX, larchwoodxylan; BeeWX, beechwoodxylan; WFA, wheat flour arabinoxylan; BirWX, birchwoodxylan.

a Mw, molecular weight presented as Mw by SDS-PAGE (Mw by gel filtration/mass spectrometry)  
b Value for pNP-Af.

## 6. Discussion

Abf51A was the only AFase identified from the genome of *C. lactoaceticus*, and showed high sequence identity with family 51 AFases from *Caldicellulosiruptor* sp. Correspondingly, the model built for Abf51A displayed a typical GH51 two-domain structure, organized with a (β/α)<sub>8</sub> folded catalytic domain and a β-sandwich folded domain at the C-terminal. The conserved Glu173 and Glu292 were proposed to be the acid-base and nucleophile catalytic residue, respectively. Additionally, like Abf51A, some GH51 AFases also had hexameric structure in solution [15-17]. All these results were in agreement with crystal structures of GH51 AFase from *G. stearothermophilus* T-6 [PDB: 1PZ3] [18], *C. thermocellum* [PDB: 2C7F] [19], *T. maritima* MSB8

[PDB: 4ATW] [20], *T. petrophila* RKU-1 [PDB: 3S2C] [21] and *T. xylanilyticus* [PDB: 2VRK] [22]. So far, GH51 Tm-AFase from *T. thermarum* DSM5069 was the most thermophilic AFase purified and characterized with optimal activity up to 95°C [23]. Recombinant Abf51A also showed excellent thermostability with optimal activity at 80°C, responding to the characterization of other thermophilic AFases [24-26]. Substrate specificity study revealed that Abf51A had narrow substrate range and mainly cleaved the glycosidic bond linked in *p*NP-Af. Abf51A was also active on *p*NP-β-Xyl, but in contrast to GH 43 AFases, this activity was rather low. Similar substrate preference had also been described with some GH51 enzymes [27-35]. The explanation for this flexibility depended on the stereochemical relatedness of α-L-arabinosyl and β-D-xylopyranosyl moieties [36]. However, no biofunctionality had been claimed on this basis [37]. To date, the majority of characterized GH51 members displayed only AFase activity, while only a few were confirmed to be multi-functional. For example, the THSAbf from *Paenibacillus* sp. displayed both AFase and endo-xylanase activity, and was active on 4-*O*-methylglucuronoxylans from birch and beechwood [38]. In all cases, it was worth noticing that Abf51A had significant higher activity with *p*NP-Af as substrate although the affinity was not outstanding. As for Abf51A, it also displayed expected activity towards AX and GAXs including CCX and SCX. AX, appearing as the neutral or slightly acidic polymer, consists of β-1,4-linked D-xylopyranosyl residues to which α-L-arabinofuranosyl units are linked at C(O)-2 and/or C(O)-3 position. Moreover, phenolic acids such as ferulic and coumaric acids have been found to be esterified to the C(O)-5 position of arabinose [39]. According to statistics, AX has been extracted from a variety of the main commercial cereals including wheat, barley, oat, rye, rice and corn, sorghum as well as some grasses [40,41]. Generally, AX from rye, wheat and sorghum on average has a relatively high degree of substituted xylose residues, whereas both oats and barley contain higher levels of β-glucan than AX [39]. For example, the proportion of AX was once isolated from barley endosperm, accounting for just 25% of the total polysaccharides in cell walls [40]. However, the degree of substitution referring to the arabinose moieties on the xylose backbone varies significantly according to the cereal species. OSX, as a popular representative of AX investigated in many literatures, contains α-L-arabinofuranosyl residue at the *O*-3 position of the linear xylan backbone at every eight to ten xylose molecules [41]. In earlier reports, CCX revealed the existence of two structurally different forms: a low-branched water-insoluble GAX with about 95% of the backbone unattached, and a water-soluble heteroxylan with over 15% of the backbone substituted [41,42]. While xylan extracted from corn fiber contains about 52% xylose, along with 34% arabinose [43]. In contrast, heteroxylan isolated from sugarcane bagasse is typical acetylated

GAX, with a glycosyl substitution molar ratio between the xylose units, *O*-3 linked terminal arabinose units and the *O*-2 linked glucuronic units of 10:0.5:0.1 [44]. In view of these facts, the detectable activity of Abf51A with polymeric xylan was due to the hydrolysis of the C(O)-2- and/or C(O)-3-linkages between α-L-arabinofuranosyl substituents and backbone residues of AX and these GAXs. From a larger perspective, the available fermentable sugars are essential for efficient conversion of lignocellulosic biomass to ethanol and other valuable bio-based products. Hence the importance of side-chain-degrading enzymes like AFase comes from the fact that these substituents extensively exist in the cross-linking structure of hemicellulose within plant cell wall. Besides, the enzymatic products from these heteroxylan, for instance, arabinoxylan-oligosaccharides, have been testified to be physiologically functional polysaccharides that exert prebiotic activities. Synergy of hydrolytic enzymes for complete degradation of arabinose-rich hemicelluloses to basic units has been successfully recorded in previous studies, which is a feasible way to improve sugar yield at low cost. For example, novel synergistic associations among AFase, endo-β-1,4-xylanase and endo-1,4-β-mannanase had already been established to degrade the hemicelluloses of alkaline pre-treated sugarcane bagasse. Respond to the results of this study, significant synergistic effects between AFase and other xylanolytic degrading enzymes had also been widely acknowledged. In these cases, efficient processing of xylan could be achieved by the predominant de-polymerization of the main bone by endo-β-1,4-xylanase to assimilate oligosaccharides, together with branch removal by the de-branching xylanolytic system. Cooperation *C. stercorarium* ArfB with endo-xylanase on OSX yielded products of arabinose, xylose, xylobiose and xylotriose after prolonged incubation. *Paenibacillus* sp. JDR-2 assimilated methylglucuronoxylan from sugarcane by the synergy between AFase and GH10/GH67 xylan utilization enzymes. Hence, when Abf51A was incubated with endo-β-1,4-xylanase Xyn10A to simultaneously degrade AX and GAXs, more reducing sugars were released. However, the difference in the initial reaction rates and products for the tested substrates might be attributed to the distribution diversity of substituents and the linkage preference of AFase. In conclusion, the GH51 AFases Abf51A from extremely thermophilic bacterium *C. lactoaceticus* exhibited remarkable activity on both *p*NP-Af and *p*NP-β-D-Xyl. Abf51A showed optimal activity at 80°C and pH 5.5 with favorable thermostability. Moreover, Abf51A has the accessory function of L-arabinofuranosyl substituents removal and acted synergistically with endo-β-1,4-xylanase in AX and GAX degradation. The excellent thermostability, hydrolytic properties and synergism with endo-β-1,4-xylanase suggested that Abf51A was an optimum enzyme for pentoses production from arabinose-rich xyans.

## 7. Acknowledgements

This study was funded by National 863 Project (grant number 2014AA021905); Open Funding Project of the National Key Laboratory of Biochemical Engineering (grant number Y525041178); 100 Talents Program of Chinese Academy of Sciences; Strategic Biological Resources Service Network Plan of Chinese Academy of Sciences (grant number ZSYS-015).

## References

1. Arti D, Jung TY, Jang MU, Kim TJ (2012) Structural analysis of  $\alpha$ -L-arabinofuranosidase from *Thermotoga maritima* reveals characteristics for thermostability and substrate specificity. *J Microbiol Biotechnol* 22(12):1724-1730. doi:10.4014/jmb.1208.08043
2. Beukes N, Pletschke BI (2011) Effect of alkaline pre-treatment on enzyme synergy for efficient hemicellulose hydrolysis in sugarcane bagasse. *Bioresour Technol* 102(8):5207-5213. doi:10.1016/j.biortech.2011.01.090
3. Birgisson H, Fridjonsson O, Bahrani-Mougeot FK, Hreggvidsson GO, Kristjansson JK, Mattiasson B (2004) A new thermostable  $\alpha$ -L-arabinofuranosidase from a novel thermophilic bacterium. *Biotechnol Lett* 26(17):1347-1351. doi:10.1023/B:BILE.0000045631.57073.79
4. Blumer-Schuette SE, Ozdemir I, Mistry D, Lucas S, Lapidus A, Cheng J-F, Goodwin LA, Pitluck S, Land ML, Hauser LJ (2011) Complete genome sequences for the anaerobic, extremely thermophilic plant biomass-degrading bacteria *Caldicellulosiruptor hydrothermalis*, *Caldicellulosiruptor kristjanssonii*, *Caldicellulosiruptor kronotskyensis*, *Caldicellulosiruptor owensensis*, and *Caldicellulosiruptor lactoaceticus*. *J Bacteriol* 193(6):1483-1484. doi:10.1128/JB.01515-10
5. Bouraoui H, Desrousseaux ML, Ioannou E, Alvira P, Manai M, Rémond C, Dumon C, Fernandez-Fuentes N, O'Donohue MJ (2016) The GH51  $\alpha$ -L-arabinofuranosidase from *Paenibacillus* sp. THS1 is multifunctional, hydrolyzing main-chain and side-chain glycosidic bonds in heteroxylans. *Biotechnol Biofuels* 9:140. doi:10.1186/s13068-016-0550-x
6. Canakci S, Belduz AO, Saha B, Yasar A, Ayaz FA, Yayli N (2007) Purification and characterization of a highly thermostable  $\alpha$ -L-arabinofuranosidase from *Geobacillus caldoolyolyticus* TK4. *Appl Microbiol Biotechnol* 75(4):813-820. doi:10.1007/s00253-007-0884-1
7. Canakci S, Kacagan M, Inan K, Belduz AO, Saha BC (2008) Cloning, purification, and characterization of a thermostable  $\alpha$ -L-arabinofuranosidase from *Anoxybacillus kestanbolensis* AC26Sari. *Appl Microbiol Biotechnol* 81(1):61-68. doi:10.1007/s00253-008-1584-1
8. Debeche T, Cummings N, Connerton I, Debeire P, O'Donohue MJ (2000) Genetic and biochemical characterization of a highly thermostable  $\alpha$ -L-arabinofuranosidase from *Thermobacillus xylanilyticus*. *Appl Environ Microbiol* 66(4):1734-1736. doi:10.1128/AEM.66.4.1734-1736.2000
9. Dodd D, CANN IK (2009) Enzymatic deconstruction of xylan for biofuel production. *GCB Bioenergy* 1(1):2-17. doi:10.1111/j.1757-1707.2009.01004.x
10. Ebringerová A, Hromádková Z, Alfödi J, Hřibalová V (1998) The immunologically active xylan from ultrasound-treated corn cobs: Extractability, structure and properties. *Carbohydr Polym* 37(3):231-239. doi:10.1016/S0144-8617(98)00065-4
11. Ebringerová A, Hromádková Z, Heinze T (2005) Hemicellulose. In: Heinze T (ed) *Polysaccharides I: Structure, Characterization and Use*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 1-67
12. Fincher GB (1975) Morphology and chemical composition of barley endosperm cell walls. *J I Brewing* 81(2):116-122. doi:10.1002/j.2050-0416.1975.tb03672.x
13. Grootaert C, Delcour JA, Courtin CM, Broekaert WF, Verstraete W, Van de Wiele T (2007) Microbial metabolism and prebiotic potency of arabinoxylan oligosaccharides in the human intestine. *Trends Food Sci Technol* 18(2):64-71. doi:10.1016/j.tifs.2006.08.004
14. Hövel K, Shallom D, Niefind K, Belakhov V, Shoham G, Baasov T, Shoham Y, Schomburg D (2003) Crystal structure and snapshots along the reaction pathway of a family 51  $\alpha$ -L-arabinofuranosidase. *The EMBO Journal* 22(19):4922-4932. doi:10.1093/emboj/cdg494
15. Heinze T, Petzold-Welcke K, van Dam JEG (2013) Polysaccharides: Molecular and supramolecular structures. Terminology. In: Navard P (ed) *The european polysaccharide network of excellence (EPNOE): Research initiatives and results*. Springer Vienna, Vienna, pp 23-64
16. Hespell RB (1998) Extraction and characterization of hemicellulose from the corn fiber produced by corn wet-milling processes. *J Agric Food Chem* 46(7):2615-2619. doi:10.1021/jf971040y
17. Hespell RB, O'Bryan PJ (1992) Purification and characterization of an  $\alpha$ -L-arabinofuranosidase from *Butyrivibrio fibrisolvens* GS113. *Appl Environ Microbiol* 58(4):1082-1088
18. Hoffmam ZB, Oliveira LC, Cota J, Alvarez TM, Diogo JA, de Oliveira Neto M, Citadini APS, Leite VBP, Squina FM, Murakami MT, Ruller R (2013) Characterization of a hexameric exo-acting GH51  $\alpha$ -L-arabinofuranosidase from the mesophilic *Bacillus subtilis*. *Mol Biotechnol* 55(3):260-267. doi:10.1007/s12033-013-9677-1
19. Im D-H, Kimura K-i, Hayasaka F, Tanaka T, Noguchi M, Kobayashi A, Shoda S-i, Miyazaki K, Wakagi T, Fushinobu S (2012) Crystal structures of glycoside hydrolase family 51  $\alpha$ -L-arabinofuranosidase from *Thermotoga maritima*. *Biosci, Biotechnol, Biochem* 76(2):423-428. doi:10.1271/bbb.110902
20. Jia X, Mi S, Wang J, Qiao W, Peng X, Han Y (2014) Insight into glycoside hydrolases for debranched xylan degradation from extremely thermophilic bacterium *Caldicellulosiruptor lactoaceticus*. *PLoS ONE* 9(9):e106482. doi:10.1371/journal.pone.0106482



21. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227(5259):680-685
22. Lagaert S, Pollet A, Courtin CM, Volckaert G (2014)  $\beta$ -Xylosidases and  $\alpha$ -L-arabinofuranosidases: Accessory enzymes for arabinoxylan degradation. *Biotechnol Adv* 32(2):316-332. doi:10.1016/j.biotechadv.2013.11.005
23. Lee SH, Lee Y-E (2014) Cloning, expression, and characterization of a thermostable GH51  $\alpha$ -L-arabinofuranosidase from *Paenibacillus* sp. DG-22. *J Microbiol Biotechnol* 24(2):236-244. doi:10.4014/jmb.1308.08078
24. Lever M (1972) A new reaction for colorimetric determination of carbohydrates. *Anal Biochem* 47(1):273-279. doi:10.1016/0003-2697(72)90301-6
25. Lim YR, Yoon RY, Seo ES, Kim YS, Park CS, Oh DK (2010) Hydrolytic properties of a thermostable  $\alpha$ -L-arabinofuranosidase from *Caldicellulosiruptor saccharolyticus*. *J Appl Microbiol* 109(4):1188-1197. doi:10.1111/j.1365-2672.2010.04744.x
26. Miyazaki K (2005) Hyperthermophilic  $\alpha$ -L-arabinofuranosidase from *Thermotoga maritima* MSB8: Molecular cloning, gene expression, and characterization of the recombinant protein. *Extremophiles* 9(5):399-406. doi:10.1007/s00792-005-0455-2
27. Mladenovska Z, Mathrani IM, Ahring BK (1995) Isolation and characterization of *Caldicellulosiruptor lactoaceticus* sp. nov., an extremely thermophilic, cellulolytic, anaerobic bacterium. *Arch Microbiol* 163(3):223-230. doi:10.1007/BF00305357
28. Morais de Carvalho D, Martínez-Abad A, Evtuguin DV, Colodette JL, Lindström ME, Vilaplana F, Sevastyanova O (2017) Isolation and characterization of acetylated glucuronarabinoxylan from sugarcane bagasse and straw. *Carbohydr Polym* 156:223-234 doi:10.1016/j.carbpol.2016.09.022
29. Numan MT, Bhosle NB (2006)  $\alpha$ -L-Arabinofuranosidases: The potential applications in biotechnology. *J Ind Microbiol Biotechnol* 33(4):247-260. doi:10.1007/s10295-005-0072-1
30. Paës G, Skov LK, O'Donohue MJ, Rémond C, Kastrup JS, Gajhede M, Mirza O (2008) The structure of the complex between a branched pentasaccharide and *Thermobacillus xylanilyticus* GH-51 arabinofuranosidase reveals xylan-binding determinants and induced fit. *Biochemistry* 47(28):7441-7451. doi:10.1021/bi800424e
31. Saha BC (2000)  $\alpha$ -L-Arabinofuranosidases: Biochemistry, molecular biology and application in biotechnology. *Biotechnol Adv* 18(5):403-423. doi:10.1016/S0734-9750(00)00044-6
32. Saulnier L, Sado P-E, Branlard G, Charmet G, Guillon F (2007) Wheat arabinoxylans: Exploiting variation in amount and composition to develop enhanced varieties. *J Cereal Sci* 46(3):261-281. doi:10.1016/j.jcs.2007.06.014
33. Sawhney N, Preston JF (2014) GH51 arabinofuranosidase and its role in the methylglucuronarabinoxylan utilization system in *Paenibacillus* sp. JDR-2. *Appl Environ Microbiol* 80(19):6114-6125.doi:10.1128/AEM.01684-14
34. Schwarz WH, Bronnenmeier K, Krause B, Lottspeich F, Staudenbauer WL (1995) Debranching of arabinoxylan: Properties of the thermoactive recombinant  $\alpha$ -L-arabinofuranosidase from *Clostridium stercoararium* (ArfB). *Appl Microbiol Biotechnol* 43(5):856-860. doi:10.1007/bf02431919
35. Shallom D, Belakhov V, Solomon D, Gilead-Gropper S, Baasov T, Shoham G, Shoham Y (2002) The identification of the acid-base catalyst of  $\alpha$ -arabinofuranosidase from *Geobacillus stearothermophilus* T-6, a family 51 glycoside hydrolase. *FEBS Lett* 514(2-3):163-167. doi:10.1016/S0014-5793(02)02343-8
36. Souza TACB, Santos CR, Souza AR, Oldiges DP, Ruller R, Prade RA, Squina FM, Murakami MT (2011) Structure of a novel thermostable GH51  $\alpha$ -L-arabinofuranosidase from *Thermotoga petrophila* RKU-1. *Protein Sci* 20(9):1632-1637. doi:10.1002/pro.693
37. Taylor E, Smith N, Turkenburg J, D'souza S, Gilbert H, Davies G (2006) Structural insight into the ligand specificity of a thermostable family 51 arabinofuranosidase, Araf51, from *Clostridium thermocellum*. *Biochem J* 395:31-37.doi:10.1042/BJ20051780
38. Van Dyk J, Pletschke B (2012) A review of lignocellulose bioconversion using enzymatic hydrolysis and synergistic cooperation between enzymes-factors affecting enzymes, conversion and synergy. *Biotechnol Adv* 30(6):1458-1480.doi:10.1016/j.biotechadv.2012.03.002
39. Xie J, Zhao D, Zhao L, Pei J, Xiao W, Ding G, Wang Z, Xu J (2016) Characterization of a novel arabinose-tolerant  $\alpha$ -L-arabinofuranosidase with high ginsenoside Rc to ginsenoside Rd bioconversion productivity. *J Appl Microbiol* 120(3):647-660.doi:10.1111/jam.13040
40. Yan Q, Tang L, Yang S, Zhou P, Zhang S, Jiang Z (2012) Purification and characterization of a novel thermostable  $\alpha$ -L-arabinofuranosidase ( $\alpha$ -L-AFase) from *Chaetomium* sp. *Process Biochem* 47(3):472-478. doi:10.1016/j.procbio.2011.12.003
41. Yang W, Bai Y, Yang P, Luo H, Huang H, Meng K, Shi P, Wang Y, Yao B (2015) A novel bifunctional GH51 exo- $\alpha$ -L-arabinofuranosidase/endo-xylanase from *Alicyclobacillus* sp. A4 with significant biomass-degrading capacity. *Biotechnol Biofuels* 8:197.doi:10.1186/s13068-015-0366-0
42. Yang Y, Sun J, Wu J, Zhang L, Du L, Matsukawa S, Xie J, Wei DZ (2016) Characterization of a novel  $\alpha$ -L-arabinofuranosidase from *Ruminococcus albus* 7 and rational design for its thermostability. *J Agric Food Chem* 64:7546-7554.doi:10.1021/acs.jafc.6b02482
43. Yeoman CJ, Han Y, Dodd D, Schroeder CM, Mackie RI, Cann IK (2010) Thermostable enzymes as biocatalysts in the biofuel industry. *Adv Appl Microbiol* 70:1-55.doi:10.1016/S0065-2164(10)70001-0

44. Zhang S, Li W, Smith CJ, Musa H (2015) Cereal-derived arabinoxylans as biological response modifiers: Extraction, molecular features, and immune-stimulating properties. *Crit Rev Food Sci Nutr* 55(8):1035-1052.[doi:10.1080/10408398.2012.705188](https://doi.org/10.1080/10408398.2012.705188)